
TECHNICAL DATA SHEET

Epimedium Extract

Catalog No.: NST-8-4

Extract Profile

Botanical source	<i>E. brevicornum Maxim.</i>
Part used	Leaf
Standardization	Icariin $\geq 10.0\%$
Extraction solvent(s)	Water and ethanol
Extract ratio / DER	20:1
Composition	90-100% Epimedium Extract; 0-10% Maltodextrin
Carrier / excipient	Maltodextrin (0-10%)

Applications

Suitable for cosmetic formulations and food supplement applications.

Solubility & Dispersibility

Water	Slightly soluble to dispersible
Oil	Insoluble

Cosmetic Formulation Guide

Recommended use level	0.1-1.0%
------------------------------	----------

Food Supplement Application Guide

Suitable dosage forms	Hard capsules (vegetarian or gelatin); tablets
Typical recommended use level	200-500 mg/day

FORMULATION & STABILITY GUIDANCE

Most stable under mildly acidic to neutral conditions (pH 4.0-7.0). Strong acids or strong bases may cause hydrolysis of the glycosidic bonds in icariin.

Relatively stable under normal processing temperatures, but prolonged exposure to temperatures above 80 °C should be avoided. Protect from direct light and moisture.

Product quality specifications, analytical methods, storage conditions and shelf life are provided in the current Quality Specification (QS) for this catalog item.

Regulatory Disclaimer

The recommended use levels provided in this Technical Data Sheet are intended for general technical guidance only and do not constitute regulatory approval or legal advice. Applicable requirements, restrictions, and permitted use levels may vary depending on the product category and country of sale. It is the responsibility of the manufacturer or formulator of the finished product to verify compliance with all applicable national and regional regulations in the intended market.